

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/269183640>

Low-level Laser (Light) Therapy Increases Mitochondrial Membrane Potential and ATP Synthesis in C2C12 Myotubes with a Peak Response at 3–6 h

Article in *Photochemistry and Photobiology* · December 2014

DOI: 10.1111/php.12397 · Source: PubMed

CITATIONS

50

READS

1,188

8 authors, including:



Cleber Ferraresi
Brasil University

54 PUBLICATIONS 681 CITATIONS

[SEE PROFILE](#)



Ying-Ying Huang
Massachusetts General Hospital

132 PUBLICATIONS 6,804 CITATIONS

[SEE PROFILE](#)



Marcelo Sousa
Bright Photomedicine, Brazil

13 PUBLICATIONS 177 CITATIONS

[SEE PROFILE](#)



Vanderlei Bagnato
University of São Paulo

313 PUBLICATIONS 2,312 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Electrophysical agents effects [View project](#)



Transcranial Photobiomodulation for Treatments of Psychiatric Disorders in Animal Models [View project](#)

Low-level Laser (Light) Therapy Increases Mitochondrial Membrane Potential and ATP Synthesis in C2C12 Myotubes with a Peak Response at 3–6 h

Cleber Ferraresi^{1,2,3,4}, Beatriz Kaippert^{4,5}, Pinar Avci^{4,6}, Ying-Ying Huang^{4,6}, Marcelo V. P. de Sousa^{4,7}, Vanderlei S. Bagnato^{2,3}, Nivaldo A. Parizotto^{1,2} and Michael R. Hamblin^{4,6,8}

¹Laboratory of Electrothermophototherapy, Department of Physical Therapy, Federal University of Sao Carlos, Sao Carlos, SP, Brazil

²Post-Graduation Program in Biotechnology, Federal University of Sao Carlos, Sao Carlos, SP, Brazil

³Optics Group, Physics Institute of Sao Carlos, University of São Paulo, Sao Carlos, SP, Brazil

⁴Wellman Center for Photomedicine, Massachusetts General Hospital, Boston, MA

⁵Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

⁶Department of Dermatology, Harvard Medical School, Boston, MA

⁷Laboratory of Radiation Dosimetry and Medical Physics, Institute of Physics, Sao Paulo University, Sao Carlos, SP, Brazil

⁸Harvard-MIT Division of Health Science and Technology, Cambridge, MA

Received 2 September 2014, accepted 25 November 2014, DOI: 10.1111/php.12397

ABSTRACT

Low-level laser (light) therapy has been used before exercise to increase muscle performance in both experimental animals and in humans. However, uncertainty exists concerning the optimum time to apply the light before exercise. The mechanism of action is thought to be stimulation of mitochondrial respiration in muscles, and to increase adenosine triphosphate (ATP) needed to perform exercise. The goal of this study was to investigate the time course of the increases in mitochondrial membrane potential (MMP) and ATP in myotubes formed from C2C12 mouse muscle cells and exposed to light-emitting diode therapy (LEDT). LEDT employed a cluster of LEDs with 20 red (630 ± 10 nm, 25 mW) and 20 near-infrared (850 ± 10 nm, 50 mW) delivering 28 mW cm^2 for 90 s (2.5 J cm^2) with analysis at 5 min, 3 h, 6 h and 24 h post-LEDT. LEDT-6 h had the highest MMP, followed by LEDT-3 h, LEDT-24 h, LEDT-5 min and Control with significant differences. The same order (6 h > 3 h > 24 h > 5 min > Control) was found for ATP with significant differences. A good correlation was found ($r = 0.89$) between MMP and ATP. These data suggest an optimum time window of 3–6 h for LEDT stimulate muscle cells.

INTRODUCTION

Mitochondria are the organelles responsible for energy production in cells and for this reason have a very important role in cellular function and maintenance of homeostasis. This organelle has an intriguing and well-designed architecture to generate adenosine triphosphate (ATP) that is the basic energy supply for all cellular activity (1,2).

Mitochondria contain a respiratory electron transport chain (ETC.) able to transfer electrons through complexes I, II, III and

IV by carrying out various redox reactions in conjunction with pumping hydrogen ions (H^+) from the mitochondrial matrix to the intermembrane space. These processes generate water as the metabolic end-product, as oxygen is the final acceptor of electrons from the ETC., that is coupled with synthesis of ATP when H^+ ions return back into mitochondrial matrix through complex V (ATP synthase), thus completing the ETC. Changes in the flow of electrons through the ETC. and consequently in H^+ pumping produce significant modulations in the total proton motive force and ATP synthesis. These changes can be measured by mitochondrial membrane potential (MMP) and content of ATP (1).

Since the earliest evidence that low-level laser (light) therapy (LLLT) can increase ATP synthesis (3,4), several mechanisms of action have been proposed to explain LLLT effects on mitochondria. One of the first studies reported increased MMP and ATP synthesis measured at an interval of 3 min after LLLT (3). Years later, other authors extended the measurement of this “extra” ATP-induced by LLLT in HeLa (human cervical cancer) cells (4). With intervals of 5 to 45 min, these authors found no change in ATP synthesis during the first 15 min after LLLT, but after 20–25 min ATP levels increased sharply and then came back to control levels at 45 min (4).

More recent studies have reported LLLT effects on mitochondria in different types of cells (5–9). In neural cells LLLT seems to also increase MMP, protect against oxidative stress (5) and increase ATP synthesis in intact cells (without stressor agents) (6). In mitochondria from fibroblast cells without stressor agents, LLLT also increased ATP synthesis and mitochondrial complex IV activity in a dose-dependent manner (7). In myotubes from C2C12 cells, LLLT could modulate the production of reactive oxygen species (ROS) and mitochondrial function in a dose-dependent manner in intact cells or in cells stressed by electrical stimulation (9).

Increases in mitochondrial metabolism and ATP synthesis have been proposed by several authors as a hypothesis to explain

*Corresponding author email: hamblin@helix.mgh.harvard.edu (Michael R. Hamblin)

© 2014 The American Society of Photobiology

LLLT effects on muscle performance when used for muscular preconditioning or muscle recovery postexercise (10–12). However, there is a lack in the literature to identify immediate and long-term effects of LLLT on mitochondrial metabolism and ATP synthesis in skeletal muscle cells that in turn could confirm these hypotheses.

This study aimed to identify the time-response for LLLT by light-emitting diode therapy (LEDT) in modulation of MMP and ATP content in myotubes from C2C12 intact cells (mouse muscle cells) only under the stress of the culture. Moreover, the second objective was to correlate MMP with ATP content within a time range of 5 min to 24 h after LLLT. Our goal was to find the best time-response for LLLT which could be useful in future experimental and clinical studies investigating muscular preconditioning, muscle recovery postexercise or any other photomodulation in muscle tissue.

MATERIALS AND METHODS

Cell culture. C2C12 cells were kindly provided by the Cardiovascular Division of the Beth Israel Deaconess Medical Center, Harvard Medical School, USA. Cells were grown in culture medium (DMEM, Dulbecco's Modified Eagle's Medium - Sigma-Aldrich) with fetal bovine serum (20% FBS - Sigma-Aldrich) and 1% antibiotic (penicillin and streptomycin) in humidified incubator at 37°C and 5% CO₂.

C2C12 cells were cultured and a total of 1.71×10^5 cells approximately were counted in a Neubauer chamber. Next, these cells were distributed equally into 30 wells (approximately 5.7×10^3 cells per well) into two different plates:

- 15 wells in black plate (Costar[®] 96-Well Black Clear-Bottom Plates) for analysis of MMP.
- 15 wells in white plate (Costar[®] 96-Well White Clear-Bottom Plates) for analysis of ATP synthesis.

Moreover, both plates were subdivided into five columns with three wells per column (triplicate):

- 1 LEDT-Control: no LEDT applied to the cells.
- LEDT-5 min: LEDT applied to the cells and assessments of ATP and MMP after 5 min.
- LEDT-3 h: LEDT applied to the cells and assessments of ATP and MMP after 3 h.
- LEDT-6 h: LEDT applied to the cells and assessments of ATP and MMP after 6 h.
- LEDT-24 h: LEDT applied to the cells and assessments of ATP and MMP after 24 h.

After plating C2C12 cells were cultured for 9 days in culture medium (DMEM) containing 2% heat-inactivated horse serum (Sigma-Aldrich) in a humidified incubator at 37°C and 5% CO₂ to induce cell differentiation into myotubes, as described in a previous study (9). At the 10th day, LEDT-24 h group received LEDT. At 11th day all remaining groups received LEDT and were assessed for ATP and MMP at specific times in accordance with each group.

Light-emitting diode therapy (LEDT). A cluster of 40 LEDs (20 red – 630 ± 10 nm; 20 infrared – 850 ± 20 nm) with a diameter of 76 mm was used in this study. The cluster was positioned at a distance of 156 mm from the top of each plate and irradiation lasted 90 s with fixed parameters as described in Table 1. Each group of wells received LEDT individually, and all others wells of each plate (groups) were covered with aluminum foil to avoid light irradiation (Fig. 1). LEDT parameters were measured and calibrated using an optical energy meter PM100D Thorlabs[®] and sensor S142C (area of 1.13 cm²). In addition, we chose use red and near-infrared light therapy at the same time to promote a double band of absorption by cytochrome c oxidase (Cox) based on specific bands of absorption reported previously (2,13–16). The room temperature was controlled (22–23°C) during LEDT irradiation, which did not increase temperature on the top of plates more than 0.5°C. This increase of 0.5°C was dissipated to room within 2 min after LEDT.

Mitochondrial membrane potential (TMRM) assay. This analysis was performed using cells placed into black plate. MMP was assessed using tetramethyl rhodamine methyl ester (TMRM – Invitrogen/Molecular Probes) at a final concentration of 25 nM. Nuclei of myotubes from

Table 1. All parameters of light-emitting diode therapy (LEDT). Control did not receive LEDT.

Number of LEDs (cluster): 40 (20 infrared-IR and 20 red-RED)
Wavelength: 850 ± 20 nm (IR) and 630 ± 10 nm (RED)
LED spot size: 0.2 cm ²
Pulse frequency: continuous
Optical output of each LED: 50 mW (IR) and 25 mW (RED)
Optical output (cluster): 1000 mW (IR) and 500 mW (RED)
LED cluster size: 45 cm ²
Power density (at the top of plate): 28 mW cm ²
Treatment time: 90 s
Cluster energy density applied on the top plate: 2.5 J cm ²
Application mode: without contact
Distance from plate or power meter: 156 mm




Figure 1. Myotubes from C2C12 cells. Experimental setup for irradiation of the white and black plates containing myotubes from C2C12 cells using light-emitting diode therapy (LEDT) without contact.

C2C12 cells were labeled using Hoechst (Sigma-Aldrich) at a concentration of 1 mg mL⁻¹. Each well was incubated for 30 min, 37°C and 5% CO₂ with 100 μL of solution containing TMRM and Hoechst. Next, this solution was carefully removed from each well and added 100 μL of buffer solution containing HBSS (Hank's Balanced Salt Solution – Life Technologies Corporation) and 15 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid – Life Technologies Corporation). The myotubes were imaged in a confocal microscope (Olympus America Inc. Center Valley, PA) using an excitation at 559 nm and emission at 610 nm. Three random fields per well were imaged with a magnification of 40× water immersion lens. Images were exported and TMRM fluorescence incorporation into mitochondrial matrix was measured using software Image J (NIH, Bethesda, MD).

Adenosine triphosphate (ATP) assay. This analysis was performed using cells placed into white plate. First, the medium was carefully removed from each well followed by addition of 50 μL per well of CellTiter Glo Luminescent Cell Viability Assay reagent (Promega). After 10 min of incubation at room temperature (25°C), luminescence signals were measured in a SpectraMax M5 Multi-Mode Microplate Reader (Molecular Devices, Sunnyvale, CA) with integration time of 5 s to increase low signals (17). A standard curve was prepared using ATP standard (Sigma) according to manufacturer's guideline and then ATP concentration was calculated in nanomol (nmol) per well.

Pearson product-moment correlation coefficient (Pearson's r). The correlation between TMRM and ATP content in myotubes from C2C12 cells was calculated using Pearson's r. The r values were interpreted as recommended previously (18): 0.00–0.19 = none to slight; 0.20–0.39 = low; 0.40–0.69 = modest; 0.70–0.89 = high; and 0.90–1.00 = very high.

Sample size calculation. The sample size was calculated based on that necessary to obtain significant differences among all groups with ATP

content. The statistical power of 80% and the effect size (greater than 0.75) were found to be satisfactory.

Statistical analysis. Shapiro–Wilk’s *W* test verified the normality of the data distribution. ATP and TMRM were compared among all groups using one-way analysis of variance (ANOVA) with Tukey HSD post hoc test. Pearson product-moment correlation coefficient (Pearson’s *r*) was conducted between TMRM and ATP. Significance was set at $P < 0.05$.

RESULTS

Mitochondrial membrane potential (TMRM)

LEDT-6 h group increased MMP (10.77 AU, SEM 0.88) compared to: Control (3.79 AU, SEM 0.46): $P < 0.001$; LEDT-5 min (4.11 AU, SEM 0.52): $P < 0.001$; LEDT-24 h (4.91 AU, SEM 0.47): $P = 0.001$. LEDT-3 h (7.87 AU, SEM 0.59) increased MMP compared to Control ($P = 0.019$) and LEDT-5 min ($P = 0.031$). These results are graphically presented in Fig. 2. All nonsignificant results were Control *versus* LEDT-5 min ($P = 0.997$) and *versus* LEDT-24 h ($P = 0.816$); LEDT-5 min *versus* LEDT-24 h ($P = 0.935$); LEDT-3 h *versus* LEDT-6 h ($P = 0.113$) and *versus* LEDT-24 h ($P = 0.103$).

ATP assay

LEDT-6 h increased ATP contents (4.53 nmol per well, SEM 0.19) compared to: Control (1.28 nmol per well, SEM 0.05): $P < 0.001$; LEDT-5 min (2.01 nmol per well, SEM 0.16): $P < 0.001$; LEDT-24 h (2.77 nmol per well, SEM 0.16): $P = 0.007$. LEDT-3 h increased ATP contents (3.73 nmol per well, SEM 0.17) compared to Control ($P < 0.001$) and LEDT-5 min ($P = 0.008$). LEDT-24 h increased ATP contents compared to Control ($P = 0.020$). These results are graphically presented in

Fig. 3A. All nonsignificant results were Control *versus* LEDT-5 min ($P = 0.385$); LEDT-3 h *versus* LEDT-6 h ($P = 0.299$) and *versus* LEDT-24 h ($P = 0.169$); LEDT-24 h *versus* LEDT-5 min ($P = 0.338$).

Sample size

The statistical power and the effect size regarding ATP content in all groups were calculated to ensure the minimal power of 80% and large effect size (>0.75). We used the mean ATP content of each group and the highest value of standard deviation among all groups, which was observed in LEDT-6 h. Our results demonstrate a difference between groups with a statistical power of 99%, effect size of 3.42 (very large effect) and total sample size of 10, i.e. 2 wells per group (five groups). These calculations demonstrate that our sample size was small, but adequate (3 wells per group).

Pearson product-moment correlation coefficient (Pearson’s *r*)

TMRM incorporation into mitochondrial matrix of myotubes from C2C12 cells showed a high correlation ($r = 0.89$) with ATP content ($P < 0.001$). This result is presented in Fig. 3B.

DISCUSSION

This study identified a well-defined time-response for the LEDT-mediated increase in MMP and ATP synthesis in myotubes from C2C12 cells under the stress of the cell culture. The light dose used was based on previous study that already reported benefits of LLLT on mitochondria of myotubes (9). In addition, we found a strong correlation between MMP and ATP content measured




Figure 2. TMRM. Analysis of mitochondrial membrane potential using tetramethyl rhodamine methyl ester (TMRM) stained in red. Images with a magnification of 40 \times . Abbreviations: LEDT= light-emitting diode therapy; AU = arbitrary units; C = control group; 5 min = LEDT-5 min group; 24 h = LEDT-24 h group; * = statistical significance ($P < 0.05$) using one-way analysis of variance (ANOVA).




Figure 3. ATP and Pearson's r . (A) Analysis of adenosine triphosphate (ATP) content between groups. (B) Pearson product-moment correlation coefficient (Pearson's r) between ATP and mitochondrial membrane potential using TMRM. Abbreviations: LEDT = light-emitting diode therapy; TMRM = tetramethyl rhodamine methyl ester; nmol = nanomol; AU = arbitrary units; C = control group; 5 min = LEDT-5 min group; 24 h = LEDT-24 h group; * = statistical significance ($P < 0.05$) using one-way analysis of variance (ANOVA).

during a wide range from 5 min (immediate effect) to 24 h (prolonged effect). To our knowledge this is the first study investigating the time-response for light therapy modulation of mitochondrial metabolism in conjunction with ATP synthesis in muscle cells.

C2C12 is a cell line originally isolated from dystrophic muscles of C3H mice by Yaffe and Saxel (19). In culture it rapidly differentiates into contractile myotubes (muscle fibers) especially when treated with horse serum instead of fetal bovine serum. These myotubes contain multinucleated cells that express proteins characteristic of skeletal muscle such as myosin heavy chain and creatine kinase (20).

One of first effects of LLLT reported in literature was a modulation on MMP and ATP synthesis in mitochondria isolated from rat liver (3) and in HeLa cells (4). Our results are in accordance with these previous studies, showing an increased MMP and ATP synthesis in myotubes from C2C12 cells. However, light therapy seems to produce a different time-response of MMP and ATP synthesis among different cell types. While HeLa cells showed a peak of ATP synthesis around 20 min after light therapy (4), mitochondria from liver showed an immediate increase in MMP and ATP synthesis (3). In this study, we found that muscle cells need a longer time in the range of 3 h to 6 h to show the maximum effect of light therapy and convert it into a significant increase in MMP and ATP synthesis, comprising an increase around 200% to 350% over the control values. In addition, we found that 24 h after irradiation, myotubes could still produce significantly more ATP compared to LEDT-Control while LEDT-5 min showed no significant difference.

Cytochrome c oxidase (Cox) has been reported to be the main chromophore in cells exposed to red and near-infrared light (2,15,16,21). However, although Cox activity is important in the immediate effects of photon absorption, the measurement of its activity may be insufficient to confirm whether light therapy can induce "extra" ATP synthesis. For this reason, the measurement of MMP in conjunction with ATP synthesis can provide information on how fast changes occur in the electron transport chain (ETC.), and H^+ pumping from the mitochondrial matrix to the intermembrane space, as well as how much H^+ ions are returning to the mitochondrial matrix (1). In this perspective, our results are consistent with Xu *et al.* (9) who reported no immediate effects of light therapy on MMP. Moreover, although Xu *et al.*




Figure 4. Mechanism of action of LEDT on mitochondria. (A) Mitochondria of myotubes from C2C12 cells without low-level laser therapy (LLL) or light-emitting diode therapy (LEDT). There is a normal flux of electrons (red arrow) through all complexes of electron transport chain, normal pumping of H^+ , normal synthesis of ATP and modest take up of TMRM by the mitochondrial matrix. (B) Mitochondria of myotubes from C2C12 cells 3–6 h after LEDT. There is an increased flux of electrons (ticker red arrow), increased pumping of H^+ , increased synthesis of ATP and increased take up of TMRM by the mitochondrial matrix. Abbreviations: I, II, III, IV and V = complexes of the mitochondrial electron transport chain; H^+ = proton of hydrogen; $-$ = electron of hydrogen; O_2 = oxygen; H_2O = metabolic water; Q = quinone; Cox = cytochrome c oxidase; ATP = adenosine triphosphate; TMRM = tetramethyl rhodamine methyl ester.

(9) did not assess ATP content, our results showed no significant responses for ATP increment immediately after light therapy compared to control group.

Our results for MMP in conjunction with ATP content had a high correlation (Pearson's $r = 0.89$) during the time range of 5 min to 24 h, suggesting a linear and positive dependence of ATP synthesis on the value of MMP (ETC. and H^+ pumping) in muscle cells, suggesting a new and more efficient time-response or time window for LEDT stimulate muscle cells (see Fig. 4A, B). These results are very important for muscle recovery postexercise (10,11) because they suggest a prolonged effect of light therapy on ATP synthesis necessary to repair muscle damage. In addition, muscular preconditioning using light therapy for improvement of performance before a bout of exercise (12) could possibly be optimized by application at the appropriate time. However, more studies *in vivo* and clinical trials are needed to confirm our hypotheses.

Muscular preconditioning using LLLT or LEDT have been reported as therapeutic approaches to improve muscle performance in both experimental models (22–24) and in clinical trials (12). However, although this improvement reported in the literature has been significant, some studies have not found positive results (25). Furthermore, differences between groups treated with light therapy or placebo seem to be not so large. These differences reported in experimental models varied between 80% and 150% of the values found for control groups for fatigue test induced by electrical stimulation (22–24). In clinical trials these differences varied between 5% and 57% increases in number of repetitions and maximal voluntary contraction (12). Possibly these relatively modest increases could be due to allowing insufficient time necessary for the muscle cells to convert light therapy into biological responses as identified in our study for MMP and ATP synthesis. Consequently, protocols for muscular preconditioning that have been done up to now (12,22–24), i.e. generally applying light 5 min before the exercise, may not possibly achieve the best result. On the basis of our results, we suggest to wait 3–6 h after light therapy irradiation to obtain the best increase in muscle performance in muscular preconditioning regimen, as MMP and ATP availability are important for muscle performance (26,27). Once more time, we would like to remark the needed of more studies *in vivo* and clinical trials to confirm our hypotheses. At this point, it is valuable to reference two previous studies that had a similar initiative (28,29). Hayworth *et al.* (28) found increments in Cox activity 24 h after apply LEDT over rats muscles; Albuquerque-Pontes *et al.* (29) found a time window, wavelength-dependence and dose response for Cox activity increase also after LLLT in rats muscles. Both studies used animals without any kind of stress, such as this study used cells only under the stress of the cell culture. We believe that these previous studies combined with our results are extremely valuable for the discovery and understanding of mechanisms of action of LLLT on muscle tissue, and may offer guidance on the future use of LLLT in clinical practice.

Our study was designed to test one dose of light during a time-response to show that there is time-dependency for LLLT to produce secondary responses in muscle cells. For this reason, this study used a constant dose (fluence) of light as reported in a previous study (9) as well as a constant power density. As there is a possible biphasic dose response (30,31), use of different parameters such as fluence, wavelengths or irradiance could produce different responses. In addition, red and near-infrared light

therapy was delivered at the same time to take advantage of the double bands in Cox to absorb the light (2,13–16).

CONCLUSION

This is the first study reporting the benefits of mixed red and near-infrared light therapy on MMP in conjunction with ATP synthesis in myotubes from C2C12 cells (muscle cells from mice). Moreover, a well-defined time-response was found for the increase in ATP synthesis mediated by MMP increased by light therapy in myotubes.

Our data suggest that 3–6 h could be the best time-response for light therapy to improve muscle metabolism. In addition, our results lead us to think there may be possible cumulative effects if light therapy is applied at intervals less than 24 h that may have clinical relevance when LLLT is used for muscle postexercise recovery. Finally, we believe that use of light therapy for muscular preconditioning could be optimized in future studies whether the time-response for increases in ATP and MMP found in this study are taken account.

Acknowledgements—We thank Professor Zoltan Pierre Arany and his instructor Glenn C. Rowe for the C2C12 cells and Andrea Brissette for assistance with multiple roles including purchase of reagents. Cleber Ferraresi thank FAPESP for his PhD scholarships (numbers 2010/07194-7 and 2012/05919-0). MR Hamblin was supported by US NIH grant R01AI050875.

REFERENCES

- Perry, S. W., J. P. Norman, J. Barbieri, E. B. Brown and H. A. Gelbard (2011) Mitochondrial membrane potential probes and the proton gradient: A practical usage guide. *Biotechniques* **50**, 98–115.
- Karu, T. (1999) Primary and secondary mechanisms of action of visible to near-IR radiation on cells. *J. Photochem. Photobiol., B* **49**, 1–17.
- Passarella, S., E. Casamassima, S. Molinari, D. Pastore, E. Quagliarillo, I. M. Catalano and A. Cingolani (1984) Increase of proton electrochemical potential and ATP synthesis in rat liver mitochondria irradiated in vitro by helium-neon laser. *FEBS Lett.* **175**, 95–99.
- Karu, T., L. Pyatibrat and G. Kalendo (1995) Irradiation with He-Ne laser increases ATP level in cells cultivated in vitro. *J. Photochem. Photobiol., B* **27**, 219–223.
- Giuliani, A., L. Lorenzini, M. Gallamini, A. Massella, L. Giardino and L. Calza (2009) Low infra red laser light irradiation on cultured neural cells: Effects on mitochondria and cell viability after oxidative stress. *BMC Complement. Altern. Med.* **9**, 8.
- Oron, U., S. Ilic, L. De Taboada and J. Streeter (2007) Ga-As (808 nm) laser irradiation enhances ATP production in human neuronal cells in culture. *Photomed. Laser Surg.* **25**, 180–182.
- Hourel, N. N., R. T. Masha and H. Abrahamse (2012) Low-intensity laser irradiation at 660 nm stimulates cytochrome c oxidase in stressed fibroblast cells. *Lasers Surg. Med.* **44**, 429–434.
- Masha, R. T., N. N. Hourel and H. Abrahamse (2013) Low-intensity laser irradiation at 660 nm stimulates transcription of genes involved in the electron transport chain. *Photomed. Laser Surg.* **31**, 47–53.
- Xu, X., X. Zhao, T. C. Liu and H. Pan (2008) Low-intensity laser irradiation improves the mitochondrial dysfunction of C2C12 induced by electrical stimulation. *Photomed. Laser Surg.* **26**, 197–202.
- Ferraresi, C., M. R. Hamblin and N. A. Parizotto (2012) Low-level laser (light) therapy (LLLT) on muscle tissue: Performance, fatigue and repair benefited by the power of light. *Photonics Lasers Med.* **1**, 267–286.
- Borsa, P. A., K. A. Larkin and J. M. True (2013) Does phototherapy enhance skeletal muscle contractile function and postexercise recovery? A systematic review. *J. Athl. Train.* **48**, 57–67.

12. Leal-Junior, E. C., A. A. Vanin, E. F. Miranda, P. D. de Carvalho, S. Dal Corso and J. M. Bjordal (2013) Effect of phototherapy (low-level laser therapy and light-emitting diode therapy) on exercise performance and markers of exercise recovery: A systematic review with meta-analysis. *Lasers Med. Sci.* [Epub ahead of print].
13. Karu, T. I., L. V. Pyatibrat and N. I. Afanasyeva (2004) A novel mitochondrial signaling pathway activated by visible-to-near infrared radiation. *Photochem. Photobiol.* **80**, 366–372.
14. Karu, T. I. and S. F. Kolyakov (2005) Exact action spectra for cellular responses relevant to phototherapy. *Photomed. Laser Surg.* **23**, 355–361.
15. Karu, T. I., L. V. Pyatibrat, S. F. Kolyakov and N. I. Afanasyeva (2008) Absorption measurements of cell monolayers relevant to mechanisms of laser phototherapy: Reduction or oxidation of cytochrome c oxidase under laser radiation at 632.8 nm. *Photomed. Laser Surg.* **26**, 593–599.
16. Karu, T. I. (2010) Multiple roles of cytochrome c oxidase in mammalian cells under action of red and IR-A radiation. *IUBMB Life* **62**, 607–610.
17. Khan, H. A. (2003) Bioluminometric assay of ATP in mouse brain: Determinant factors for enhanced test sensitivity. *J. Biosci.* **28**, 379–382.
18. Weber, J. and D. Lamb (1970) *Statistics and Research in Physical Education*. C. V. Mosby Co., Saint Louis.
19. Yaffe, D. and O. Saxel (1977) Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature* **270**, 725–727.
20. Tannu, N. S., V. K. Rao, R. M. Chaudhary, F. Giorgianni, A. E. Saeed, Y. Gao and R. Raghov (2004) Comparative proteomes of the proliferating C(2)C(12) myoblasts and fully differentiated myotubes reveal the complexity of the skeletal muscle differentiation program. *Mol. Cell Proteomics* **3**, 1065–1082.
21. Karu, T. I. (2008) Mitochondrial signaling in mammalian cells activated by red and near-IR radiation. *Photochem. Photobiol.* **84**, 1091–1099.
22. Lopes-Martins, R. A., R. L. Marcos, P. S. Leonardo, A. C. Jr Prianti, M. N. Muscara, F. Aimbire, L. Frigo, V. V. Iversen and J. M. Bjordal (2006) Effect of low-level laser (Ga-Al-As 655 nm) on skeletal muscle fatigue induced by electrical stimulation in rats. *J. Appl. Physiol.* (1985), **101**, 283–288.
23. Leal Junior, E. C., R. A. Lopes-Martins, P. de Almeida, L. Ramos, V. V. Iversen and J. M. Bjordal (2010) Effect of low-level laser therapy (GaAs 904 nm) in skeletal muscle fatigue and biochemical markers of muscle damage in rats. *Eur. J. Appl. Physiol.* **108**, 1083–1088.
24. Santos, L. A., R. L. Marcos, S. S. Tomazoni, A. A. Vanin, F. C. Antonialli, V. D. Grandinetti, G. M. Albuquerque-Pontes, P. R. de Paiva, R. A. Lopes-Martins, P. D. de Carvalho, J. M. Bjordal and E. C. Leal-Junior (2014) Effects of pre-irradiation of low-level laser therapy with different doses and wavelengths in skeletal muscle performance, fatigue, and skeletal muscle damage induced by tetanic contractions in rats. *Lasers Med. Sci.* [Epub ahead of print].
25. Higashi, R. H., R. L. Toma, H. T. Tucci, C. R. Pedroni, P. D. Ferreira, G. Baldini, M. C. Aveiro, A. Borghi-Silva, A. S. de Oliveira and A. C. Renno (2013) Effects of low-level laser therapy on biceps braquialis muscle fatigue in young women. *Photomed. Laser Surg.* **31**, 586–594.
26. Allen, D. G., G. D. Lamb and H. Westerblad (2008) Skeletal muscle fatigue: Cellular mechanisms. *Physiol. Rev.* **88**, 287–332.
27. Ferraresi, C., T. de Brito Oliveira, L. de Oliveira Zafalon, R. B. de Menezes Reiff, V. Baldissera, S. E. de Andrade Perez, E. Matheucci Junior and N. A. Parizotto (2011) Effects of low level laser therapy (808 nm) on physical strength training in humans. *Lasers Med. Sci.* **26**, 349–358.
28. Hayworth, C. R., J. C. Rojas, E. Padilla, G. M. Holmes, E. C. Sheridan and F. Gonzalez-Lima (2010) In vivo low-level light therapy increases cytochrome oxidase in skeletal muscle. *Photochem. Photobiol.* **86**, 673–680.
29. Albuquerque-Pontes, G. M., R. D. Vieira, S. S. Tomazoni, C. O. Caires, V. Nemeth, A. A. Vanin, L. A. Santos, H. D. Pinto, R. L. Marcos, J. M. Bjordal, P. D. de Carvalho and E. C. Leal-Junior (2014) Effect of pre-irradiation with different doses, wavelengths, and application intervals of low-level laser therapy on cytochrome c oxidase activity in intact skeletal muscle of rats. *Lasers Med. Sci.* [Epub ahead of print].
30. Huang, Y. Y., A. C. Chen, J. D. Carroll and M. R. Hamblin (2009) Biphasic dose response in low level light therapy. *Dose Response* **7**, 358–383.
31. Huang, Y. Y., S. K. Sharma, J. Carroll and M. R. Hamblin (2011) Biphasic dose response in low level light therapy - an update. *Dose Response* **9**, 602–618.